arXiv:1612.01722v1 [cond-mat.stat-mech] 6 Dec 2016

Geometric explanation of anomalous finite-size scaling in high dimensions

Jens Grimm,[1] Eren Metin Elçi,[2] Zongzheng Zhou,[1] Tim Garoni,[1] and Youjin Deng[3]

[1]*ARC Centre of Excellence for Mathematical and Statistical Frontiers (ACEMS), School of Mathematical Sciences, Monash University, Clayton, VIC 3800, Australia*
[2]*School of Mathematical Sciences, Monash University, Clayton, Victoria 3800, Australia*
[3]*Hefei National Laboratory for Physical Sciences at Microscale, Department of Modern Physics, University of Science and Technology of China, Hefei 230027, China*

(Dated: December 7, 2016)

We give an intuitive geometric explanation for the apparent breakdown of standard finite-size scaling in systems with periodic boundaries above the upper critical dimension. The Ising model and self-avoiding walk are simulated on five-dimensional hypercubic lattices with free and periodic boundary conditions, by using geometric representations and recently introduced Markov-chain Monte Carlo algorithms. We show that previously observed anomalous behaviour for correlation functions, measured on the standard Euclidean scale, can be removed by defining correlation functions on a scale which correctly accounts for windings.

Finite-size scaling (FSS) is a fundamental theory within statistical mechanics, describing the asymptotic approach to the thermodynamic limit of finite systems in the neighbourhood of a critical phase transition [1, 2]. Moreover, FSS provides a key computational tool used to estimate critical exponents from Monte Carlo data [3].

It is well-known that models of critical phenomena typically possess an upper critical dimension, d_c, above which their thermodynamic behaviour is governed by critical exponents taking simple mean-field values. However, the theory of FSS in dimensions above d_c is surprisingly subtle, and remains the subject of ongoing debate [4–11].

The majority of this debate has focused on the boundary-dependent FSS of the ferromagnetic Ising model above its upper critical dimension $d_c = 4$; particularly on the case $d = 5$. Numerical observations [12] for the magnetic susceptibility χ have established an anomalous FSS behaviour $\chi \sim L^{d/2}$ at the critical point, when using periodic boundary conditions (PBC). By contrast, standard mean field behaviour $\chi \sim L^2$ is observed [6, 8] for free boundary conditions (FBC). Moreover, with periodic boundaries, it was numerically observed [7] that the correlation between spins at distance $L/2$ scales like $L^{-d/2}$, in contrast to the standard mean-field prediction $L^{-(d-2)}$ expected for free boundaries.

Recent renormalization-group arguments attempt to explain this anomalous FSS by postulating a modified scaling of the correlation length $\xi \sim L^{\wp}$ when $d > d_c$, where $\wp := d/d_c$ [13]. Moreover, an additional exponent η_Q, related to $\wp$, was introduced to explain the anomalous large-distance behaviour of the spin-spin correlation function.

In this Letter, we argue that in order to elucidate the mechanisms underlying boundary-dependent FSS, it proves useful to consider appropriate geometric representations of the investigated systems. In particular, rather than working directly with Ising spins, we study the Ising model via its high-temperature representation [14]. In addition, we also study the analogous FSS properties for the self-avoiding walk (SAW). Both the Ising and SAW models are special cases of the $O(n)$ vector model; $n = 1$ corresponding to Ising, and $n \to 0$ to SAW. Both models have upper critical dimension $d_c = 4$.



Figure 1: Average winding number, W, for Ising and SAW models with periodic boundary conditions. The number of windings is asymptotically constant in L for $d < d_c$. Above d_c, windings proliferate with increasing L. To emphasize the universal scaling, the data for Ising and SAW were translated onto a single curve for both $d = 2$ and $d = 5$. The predicted scaling $W \sim L^{\wp-1}$ with $\wp = d/d_c$ is evident for $d > d_c$.

We argue that the apparent breakdown of standard FSS for periodic systems is a manifestation of the proliferation of windings (see below for a precise definition), absent for dimensions below d_c; see Fig. 1. This motivates introducing an alternative definition of length on the torus, different from the standard Euclidean length, which accounts for the number of such windings. We refer to this length scale, which is of order $WL \sim L^{\wp}$, as the *unwrapped length*. Our numerical results show that when correlations are measured on the scale of the unwrapped length, the apparently anomalous behaviour on

the torus disappears, and standard mean-field behaviour is recovered.

Furthermore, we provide strong numerical evidence that even when measured on the Euclidean scale, the observed anomalous behaviour of critical correlations on the torus can be explained without introducing a new critical exponent η_Q. In particular, Fig. 4a and 4b confirm the following piecewise asymptotic behaviour

$$\langle s_0 s_{\mathbf{x}} \rangle_{PBC} \sim \begin{cases} \|\mathbf{x}\|^{-(d-2)}, & \|\mathbf{x}\| \leq O\left(L^{d/[2(d-2)]}\right), \\ L^{-d/2}, & \|\mathbf{x}\| \geq O\left(L^{d/[2(d-2)]}\right), \end{cases} \quad (1)$$

as conjectured in [15]. These figures also show that the analogous quantity for SAWs obeys the same scaling.

We emphasize that even though, above d_c, the Ising and SAW models exhibit the same mean-field thermodynamic behaviour on the infinite lattice, it is not *a priori* obvious that their FSS behaviours on finite tori need agree. Indeed, the essential reason that the Ising and SAW models belong to the same universality class above d_c is that both are essentially described by simple random walks. On the torus, however, simple random walk is recurrent rather than transient, implying that the corresponding Green's function does not exist. The familiar link between SAW and Ising correlations, valid on the infinite lattice, therefore breaks down on the torus.

Nevertheless, our numerical results show that the Ising and SAW correlations do indeed display the same FSS on the torus.

Geometric representations and observables.— The zero-field Ising model is defined by the Hamiltonian $\mathcal{H} = -\sum_{ij} s_i s_j$, where $s_i \in \{-1, +1\}$ is the spin on site i of a hypercubic lattice of side length L, and the sum is over nearest neighbours. Its so-called high-temperature expansion [14] provides a natural geometric representation for the spin-spin correlation function

$$g_{\text{Ising}}(\mathbf{x}) := \langle s_0 \, s_{\mathbf{x}} \rangle = \frac{\sum_{A:\partial A=\{0,\mathbf{x}\}} z^{|A|}}{\sum_{A:\partial A=\varnothing} z^{|A|}}, \quad (2)$$

where the sums are over all bond configurations A subject to the given constraint on ∂A, where ∂A denotes the set of vertices incident to an *odd* number of occupied bonds. The bond fugacity satisfies $z = \tanh(1/T)$, where T is the Ising temperature. We simulated this geometric representation using the worm algorithm of Prokofiev and Svistunov [16].

We also considered the SAW model in the grand-canonical (variable-length) ensemble. The SAW correlation function is given by

$$g_{\text{SAW}}(\mathbf{x}) := \sum_{\omega:\ 0 \to \mathbf{x}} z^{|\omega|}, \quad (3)$$

where the sum is over all SAWs beginning at the origin and ending at site $\mathbf{x}$. In contrast to typical studies of SAWs, which are performed on the infinite lattice, we consider this model to be confined to finite subsets of the hypercubic lattice, as studied for the Ising model. We simulated this ensemble using an irreversible version of the Beretti-Sokal (B-S) algorithm [17] introduced in [18].

For convenience, we measured the Ising and SAW correlation functions only along the first coordinate axis, i.e. only at $\mathbf{x} = (x, 0, \ldots, 0)$ for $0 \leq x \leq L/2$.

In addition, we measured the winding number $\mathcal{W}$:

- For the SAW model, $\mathcal{W}$ is defined as the number of windings along the first coordinate axis.
- In the Ising model, $\mathcal{W}$ is defined as the number of windings along the first coordinate axis, in the largest cluster. Measurements were taken when $\partial A = \varnothing$.

We emphasize that $\mathcal{W}$ does not distinguish between windings in the positive or negative directions, and takes strictly non-negative values. We denote the mean value of $\mathcal{W}$ by $W := \langle \mathcal{W} \rangle$.

Furthermore, for the SAW model, we additionally measured:

- The walk length $\mathcal{N}$, and its mean N.
- The unwrapped correlation function $\tilde{g}_{\text{SAW}} : \mathbb{N} \to \mathbb{R}$

$$\tilde{g}_{\text{SAW}}(u) := \sum_{\omega \in \mathcal{S}_1 : \mathcal{U}(\omega)=u} z^{\mathcal{N}(\omega)} \quad (4)$$

  where $\mathcal{S}_1$ is the set of all SAWs on $\mathbb{Z}_L^d$ which start at the origin and end on the first coordinate axis, and where the unwrapped length $\mathcal{U}$ is defined algorithmically as follows. For $\omega \in \mathcal{S}_1$, traverse ω from the origin to its endpoint, adding +1 (−1) for each step of the walk in the positive (negative) direction along the first coordinate axis.

The unwrapped length $\mathcal{U}$ simply corresponds to the length the walk would have in the infinite lattice, if the torus were unwrapped, so that periodic images are considered distinct.

Our simulations for both models were performed on the standard hypercubic lattice, using both FBC and PBC. The Ising model was simulated at the exact infinite-volume critical point in two dimensions [20], and at the estimated location of the infinite-volume critical point $z_{c,\text{Ising,5d}} = 0.113\ 915\ 0(5)$ [6] in five dimensions. The SAW model was simulated at the estimated location of the infinite-volume critical points, $z_{c,\text{SAW,2d}} = 0.379\ 052\ 277\ 758(4)$ [21], $z_{c,\text{SAW,5d}} = 0.113\ 140\ 84(1)$[18] and $z_{c,\text{SAW,6d}} = 0.091\ 927\ 86(4)$ [23], in dimensions, 2, 5 and 6, respectively. Our fitting methodology and corresponding error estimation follow standard procedures, see e.g. [25, 26]. To estimate the exponent value for a generic observable Y we performed least-squares fits to the ansatz $Y = a_Y L^{b_Y} + c_Y$.

To estimate, $g_{\text{Ising, PBC/FBC}}$ we achieved linear system sizes up to $L = 101$. Our SAW simulations were performed in the range $21 \le L \le 201$ for PBC and $51 \le L \le 401$ for FBC, respectively. A detailed analysis of autocorrelation times can be found in [24] for the worm algorithm and in [18] for the irreversible B-S algorithm.

Boundary dependent FSS, and choosing the right scale.— We now present a scaling argument which characterizes the proliferation of windings in the SAW model in terms of the exponent $\wp$. Consider a uniformly random SAW of fixed length N in $\mathbb{Z}^d$, with $d > d_c$. The second virial coefficient $B_2^{N,N}$ provides a measure of the excluded volume between a pair of such SAWs, and is believed to scale like $B_2^{N,N} \sim N^2$ (see e.g. [27]). This suggests that in order to wrap such a walk onto a torus $\mathbb{Z}_L^d$, without introducing intersections, would require $N^2 \lesssim L^d$. Considering now a variable length ensemble at z_c, we expect the mean of $\mathcal{N}$ to be of the order of its maximum, which implies $\mathbb{E}(\mathcal{N}) \sim L^{d/2}$. Figure 2 verifies this prediction. Furthermore, if one were to take a typical SAW on the torus $\mathbb{Z}_L^d$, and unwrap it into $\mathbb{Z}^d$, it would have root-mean-square displacement of order WL. But for a uniformly-random fixed-length SAW in $\mathbb{Z}^d$ with $d > d_c$, the mean-square displacement scales like the walk length. Combining this with the above observation shows that $WL \sim L^{d/4} = L^{\wp}$.

Figure 1 confirms that this prediction holds for the SAW model with $d = 5, 6$, and also for the Ising model with $d = 5$. By contrast, no proliferation of windings is observed for $d = 2$ in either model. Our fits yield $b_{W,\text{SAW,5d}} = 0.27(4)$ and $b_{W,\text{Ising,5d}} = 0.24(3)$ and $b_{W,\text{SAW, 6d}} = 0.54(7)$, in good agreement with the predicted value of $\wp - 1$.



Figure 2: FSS of the average walk length N of a critical SAW in five dimensions. As predicted in the text, on the torus we observe $N \sim L^{d/2}$. By contrast, with free boundaries we observe the standard mean-field behaviour $N \sim L^2$.

Unwrapped correlation function.— Consider simple random walk (SRW) on a finite torus $\mathbb{Z}_L^d$ with $d > 2$. If one defines $g_{\text{SRW}}(x)$ analogously to $g_{\text{SAW}}(x)$, it diverges at the SRW critical fugacity $z = 1/2d$. However, if instead one considers the unwrapped version, defined by summing over simple random walks in (4), rather than SAWs, then one immediately recovers the usual infinite-lattice simple-random walk Green's function.

This simple observation suggests that the unwrapped SAW correlation function $\tilde{g}_{\text{SAW}}(u)$ may also recover the standard mean-field behaviour. Figure 3 clearly illustrates that this is indeed the case. From the figure, we see that the unwrapped correlation function for systems with PBC displays identical scaling behaviour to the Euclidean correlation function for systems with FBC. Numerically fitting the exponent corresponding to the power-law decay $\tilde{g}_{\text{SAW,PBC}}(u) \sim u^{-b}$ yields $b = -3.02(2)$. Similarly, fitting the decay exponent for $g_{\text{SAW,FBC}}(x)$ yields $b = -3.01(2)$. Both estimates are in excellent agreement with the expected $x^{-(d-2)}$ scaling corresponding to the infinite-lattice SRW Green's function.



Figure 3: Comparison of $g_{\text{SAW, FBC}}$ and $\tilde{g}_{\text{SAW, PBC}}$. For clarity, the data for $g_{\text{SAW,FBC}}(x)$ was translated upwards.

Correlations on the Euclidean scale.— Finally, we now consider the correlation functions for the Ising and SAW models with periodic boundary conditions, on the Euclidean scale. As predicted by (1), Fig. 4a shows that two qualitatively different regions of x can be identified. At short distances, both correlation functions exhibit standard mean-field decay $x^{-(d-2)}$, while at long distances, both enter a plateau. In the inset, we fix a point in the plateau, $x = \lfloor L/4 \rfloor$, and analyze the L dependence of the correlations at this point. Our fits yield exponent values of $-2.55(12)$ and $-2.52(23)$ for SAW and Ising, respectively, in good agreement with the prediction $d/2$ from (1). This $L^{-d/2}$ scaling in the bulk is in agreement with a previous study of the Ising model presented in [7].

To further test the conjectured scaling form (1), Fig. 4b plots appropriately scaled versions of $g_{\text{SAW,PBC}}$ and $g_{\text{Ising,PBC}}$ against the dimensionless variable $y := x/L^{d/[2(d-2)]}$. The excellent data collapse provides strong evidence for the validity of (1).



Figure 4: (a) Correlation functions for SAW and Ising models with PBC, on the Euclidean scale. The inset shows FSS at the fixed point $x = \lfloor L/4 \rfloor$ leading to $g_{\text{Ising+SAW,PBC}}(\lfloor L/4 \rfloor) \sim L^{-d/2}$ for both models, in contrast to the standard mean field prediction of L^{-3}. (b) Collapse of both the Ising and SAW data with $y := x/L^{d/[2(d-2)]}$ onto the ansatz (1).

Discussion.— In this Letter, we have studied boundary-dependent FSS above d_c for both the SAW model, and for a geometric representation of the Ising model. We have established that the anomalous behaviour observed previously for the correlation functions can be explained geometrically, without any need for new critical exponents. This conclusion is in broad agreement with independent arguments made previously in [8]. In that work, an analysis of the Fourier modes of the Ising model was presented, which also refuted the need for the exponent η_Q.

Moreover, our results show that if one considers correlations of the periodic system on the scale of the unwrapped length, rather than the Euclidean length, then standard mean-field behaviour is recovered. Furthermore, the scale of the unwrapped length is shown to be governed by the exponent ♀.

Our consideration of unwrapped correlations above focused on the SAW case, largely for reasons of computational efficiency. However we expect analogous constructions to apply to the Ising case. Unlike the SAW case, there does not appear to be one unique sensible choice for the definition of unwrapped length in the Ising case. One sensible candidate would appear to be the longest path along one fixed axis between the odd-degree vertices. Unfortunately, finding the longest path in a graph is a computationally demanding task, which makes testing this conjecture challenging [28].

Finally, we note that anomalous FSS behaviour on tori has been established rigorously for percolation and the Loop Erased Random Walk (LERW). For LERW, it has been shown [19] that the mean path length scales as $L^{d/2}$, in agreement with our observations for SAW. For percolation it was conjectured that for $d > 6$ [29] the largest cluster scales as L^4 for bulk boundary conditions while it scales as $L^{2d/3}$ with periodic boundaries. This conjecture was subsequently proved, for sufficiently large dimension, for bulk boundaries in [30] and for periodic boundaries in [31]. It would be of significant interest to study such percolative questions in the general framework of the random cluster model, and examine the Ising model from this alternative geometric perspective.

Acknowledgments.— We would like to thank S. Bowly, A. Collevecchio, H. Hu, R. Kenna, M. Weigel and U. Wolff for fruitful discussions. This work was supported under the Australian Research Councils Discovery Projects funding scheme (Project Number DP140100559). It was undertaken with the assistance of resources from the National Computational Infrastructure (NCI), which is supported by the Australian Government. Furthermore, we would like to acknowledge the Monash eResearch Centre and eSolutions-Research Support Services through the use of the Monash Campus HPC Cluster. Y. Deng thanks the National Natural Science Foundation of China for their support under Grant No. 11625522 and the Fundamental Research Funds for the Central Universities under Grant No. 2340000034. J. Grimm and E. M. Elçi thank the University of Science and Technology of China for its hospitality during which this work was written.

[1] M. E. Fisher, *The theory of critical points singularities*, Proceedings of the 51st Enrico Fermi Summer School, Varenna, Italy, (1971).

[2] M. E. Fisher, M. N. Barber, *Scaling theory for finite-size effects in the critical region*, Phys. Rev. Lett. 28, 1516 (1972).

[3] M. E. J. Newman, G. T. Barkema, *Monte Carlo Methods*

in Statistical physics, Oxford University Press, (1999).

[4] P. H. Lundow, K. Markström, Non-vanishing boundary effects and quasi-first- order phase transitions in high dimensional Ising models, Nucl. Phys. B 845, 120-139 (2011).

[5] B. Berche, R. Kenna, J.-C. Walter, *Hyperscaling above the upper critical dimension*, Nuclear Physics B 865 115-132 (2012).

[6] P. H. Lundow, K. Markström, *Finite size scaling of the 5D Ising model with free boundary conditions*, Nucl. Phys. B, vol 889, p249 (2014).

[7] R. Kenna, B. Berche, *Fisher's scaling relation above the upper critical dimension*, EPL, 105 26005 (2014).

[8] M. Wittmann, A. P. Young, *Finite-size scaling above the upper critical dimension*, Phys. Rev. E 90, 062137 (2014).

[9] E. Flores-Sola, B. Berche, R. Kenna, M. Weigel, *Role of Fourier Modes in Finite-Size Scaling above the Upper Critical Dimension*, Phys. Rev. Lett. 116, 115701 (2016).

[10] R. Kenna, B. Berche, *Universal Finite-Size Scaling for Percolation Theory in High Dimensions*, arXiv:1606.00315 [cond-mat.stat-mech] (2016).

[11] P. H. Lundow, K. Markström, *The scaling window of the 5D Ising model with free boundary conditions*, arXiv:1601.04053 [cond-mat.stat-mech] (2016).

[12] E. Luijten, K. Binder, H.W.J. Blöte, *Finite-size scaling above the upper critical dimension revisited: the case of the five-dimensional Ising model*, Eur. Phys. J. B 9, 289297, (1999).

[13] R. Kenna, B. Berche, *Scaling and Finite-Size Scaling above the Upper Critical Dimension*, Order, Disorder and Criticality: Advanced Problems of Phase Transition Theory (2015).

[14] C. J. Thompson, *Mathematical Statistical Mechanics*, Princeton University Press (1979).

[15] V. Papathanakos, *Finite-Size Effects in High-Dimensional Statistical Mechanical Systems: The Ising Model With Periodic Boundary Conditions*, PhD Thesis, Princeton University (2006).

[16] N.Prokof'ev, B. Svistunov, *Worm Algorithms for Classical Statistical Models*, Phys. Rev. Lett. 87, 160601 (2001).

[17] A. Berretti, A. D. Sokal, *New Monte Carlo method for the self- avoiding walk*, J. Stat. Phys. 40, 483 (1985).

[18] H. Hu, X. Chen, Y. Deng, *Irreversible Markov chain Monte Carlo algorithm for the self-avoiding walk*, arXiv:1602.01671 [cond-mat.stat-mech]

[19] I. Benjamini, G. Kozma, *Loop-Erased Random Walk on a Torus in Dimensions 4 and Above*, Commun. Math. Phys. 259, 257-286 (2005).

[20] R. J. Baxter, Exactly solved models in statistical mechanics, London: Academic Press Inc. [Harcourt Brace Jovanovich Publishers] (1982).

[21] I. Jensen, *A parallel algorithm for the enumeration of self- avoiding polygons on the square lattice*, J. Phys. A: Math. Gen. 36, 5731 (2003).

[22] H.-P. Hsu, P. Grassberger, *Polymers confined between two par- allel plane walls*, J. Chem. Phys. 120, 2034 (2004).

[23] A. L. Owczarek, T. Prellberg, *Scaling of self-avoiding walks in high dimensions*, J. Phys. A: Math. Gen. 34, 5773 (2001).

[24] Y. Deng, T. M. Garoni, A. D. Sokal, *Dynamic Critical Behavior of the Worm Algorithm for the Ising Model*, Phys. Rev. Lett. 99, 110601 (2007).

[25] P. Young, *Everything You Wanted to Know About Data Analysis and Fitting but Were Afraid to Ask*, SpringerBriefs (2015).

[26] A. D. Sokal, *Monte Carlo Methods in Statistical Mechanics: Foundations and New Algorithms*, Lectures at the Cargèse Summer School on 'Functional Integration: Basics and Applications' (1996).

[27] B. Li, N. Madras, A. D. Sokal, *Critical exponents, hyperscaling, and universal amplitude ratios for two-and three-dimensional self-avoiding walks*, Journal of Statistical Physics 80.3-4, 661-754 (1995).

[28] A. Schrijver, *Combinatorial Optimization: Polyhedra and Efficiency, Volume 1, Algorithms and Combinatorics*, 24, Springer, (2003).

[29] M. Aizenman, *On the Number of Incipient Spanning Clusters*, Nucl. Phys. B 485 [FS], 551-582 (1997).

[30] T. Hara, *Decay of correlations in nearest-neighbor self-avoiding walk, percolation, lattice trees and animals.*, Ann. Probab. Volume 36, Number 2, 530-593, (2008).

[31] M. Heydenreich, R. van der Hofstad, *Random Graph Asymptotics on High-Dimensional Tori*, Commun. Math. Phys. 270 335 (2007).